



ImmuCast

Release Notes

v5.24.0



Support Services

For general support on this product, contact your system administrator or help desk. For up-to-date documentation, visit the STC Documentation Portal at <https://documentation.stchome.com/>.

Connect with Us on Social Media



Copyrights and Trademarks

© 2019 by Scientific Technologies Corporation (STC). All rights reserved.

This documentation describes the following: ImmuCast 5.24.0 (and IWeb Forecaster) release notes

No part of this publication may be altered, reproduced, transmitted, stored in a retrieval system, or translated into any human or computer language, in any form or by any means, electronic, mechanical, magnetic, optical, or otherwise, without the prior written permission of the copyright owner.

This document, along with any embedded media, is provided for informational purposes only. Scientific Technologies Corporation (STC) provides this material "as is" and does not assert that this document is error free. The information contained herein may not necessarily accurately represent the current or planned functions of the application and may be subject to significant and frequent modification. Scientific Technologies Corporation (STC) may change this document, any embedded content, and the product described herein at any time. Any changes will be incorporated in new versions of this document.

Scientific Technologies Corporation (STC) recognizes the rights of the holders of all trademarks used in its publications.

This document may provide hyperlinks to third-party websites or access to third-party content. Links and access to third-party sites are provided for your convenience only. Scientific Technologies Corporation does not control, endorse, or guarantee third-party content and is not responsible for any content, associated links, resources or services associated with a third-party site. Scientific Technologies Corporation shall also not be liable for any loss or damage associated with your use of any third-party content. (20181212)

Table of Contents

Introduction.....	1
Apply the Release.....	1
For ImmuCast (Stand-Alone Forecaster).....	1
Patient Groups Affected by this Release (v5.24.0)	1
New Features/Improvements	2
Bug Fix Summary	2
Details for v5.24.0.....	4
Known Issues for v5.24.0	6
HPV	6
Hib	6
Pneumococcal	7
Rotavirus.....	8
Zostavax	9
Planned Logic Changes.....	9
Product Documentation	10

Introduction

This release contains schedule changes that may affect the group of patients you select to be re-forecasted. Detailed descriptions and test cases follow below in the ticket descriptions.

Also note that Forecast, Forecaster, and ImmuCast are used interchangeably throughout this document.

Apply the Release

Apply the release by executing either the included `forecast.bat` or `forecast.sh` file. Prior forecast releases through version 5.23.0 should have already been applied. Log files, which are created in the same folder from where the release is executed, can be reviewed for errors. To determine the current version of ImmuCast, execute this statement from SQLPlus:

```
SQL> select max(version) from h33_forecast_version where insert_stamp =  
(select max(insert_stamp) from h33_forecast_version);
```

If the version number returned is not 5.23.0 or higher, download and apply the previous releases prior to applying this release. Log files are created in the folder from where the release is executed and can be reviewed for errors.

For ImmuCast (Stand-Alone Forecaster)

After applying the release, restart Tomcat to enable and cache any new vaccine codes into memory.

Patient Groups Affected by this Release (v5.24.0)

This release updates vaccine codes and influenza forecasting for 2020-21. Patients that were administered vaccine that corresponds to the new 2020 CVX codes for influenza would not have been forecasted prior to this release.

New Features/Improvements

The following are new features, functionalities, or enhancements in this version of ImmuCast.

CVX Codes Updates

CVX	CPT	CDC Short Description	CDC Long Description
203	90619	meningococcal polysaccharide (groups A, C, Y, W-135) TT conjugate	meningococcal polysaccharide (groups A, C, Y, W-135) tetanus toxoid conjugate vaccine .5mL dose, preservative free
205	90694	Influenza vaccine, quadrivalent, adjuvanted	influenza, seasonal vaccine, quadrivalent, adjuvanted, .5mL dose, preservative free
206		Smallpox monkeypox vaccine (National Stockpile)	smallpox monkeypox vaccine, live attenuated, preservative free (National Stockpile)

Bug Fix Summary

Ticket	Description
HDSD-1803	Resolved the issue where Influenza Dose 2 was marked as invalid when a child turns 9 in same flu season.
HDSD-1515	Resolved the issue where STC Test Case Flu 2019-3 Shows second flu in second season as Invalid.
HDSD-2521	Updated the short description to display as "Influenza vaccine, quadrivalent, adjuvanted".
HDSD-2594	The new CVX codes for influenza have been added to the forecast logic (no test cases included).
	2020-21 ACIP Influenza Recommendations – no changes from prior year needed. Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices – United States, 2020-21 Influenza Season

Details for v5.24.0

The following lists the detailed information about each of the tickets addressed in v5.24.0. Please note: When multiple service desk tickets address the same issue, the test cases below may not include all scenarios from all tickets.

Key	Vaccine Group	Description						
		Refer to Bug Fix Summary above						
Test Scenario		Antigen	Vacc Date	Valid	Rec Date	Min Date	Past Due	Comment
HDSD-1803 Patient receives dose 1 influenza at 8 years of age and turns 9 years old during same flu season.	DOB: 01/07/2011	Influenza CVX 161	11/07/2019	Y	12/05/2019	12/05/2019	01/01/2020	
		Influenza CVX 161	02/29/2020	Y	07/01/2020	07/01/2020	07/28/2020	Initial 2 dose influenza complete.
Test Scenario		Antigen	Vacc Date	Valid	Rec Date	Min Date	Past Due	Comment
HDSD-1515 Scenario 1: Patient receives dose 1 influenza in a prior season. Two doses are administered the next season. Dose 2 shows as Inadvertent but should be Valid.	DOB: 01/01/2018	Influenza CVX 161	07/01/2018	Y	07/29/2018	07/29/2018	08/25/2018	
		Influenza CVX 161	10/01/2019 (Eval Date: 10/01/2019)	Y	10/29/2019	10/29/2019	11/25/2019	
		Influenza CVX 161	11/01/2019	Y	07/01/2020	07/01/2020	07/28/2020	

Test Scenario		Antigen	Vacc Date	Valid	Rec Date	Min Date	Past Due	Comment
Scenario 2: Doses 1 and 2 same as Scenario 1 but with evaluation date of 09/10/2020 and no Dose 3. For 2020-21 influenza season, patient has 2 prior doses and is forecast for current season.	DOB: 01/01/2018	Influenza CVX 161	07/01/2018	Y	07/29/2018	07/29/2018	08/25/2018	
		Influenza CVX 161	10/01/2019 (Eval Date: 09/10/2020)	Y	07/01/2020	07/01/2020	07/28/2020	

Known Issues for v5.24.0

The following are known issues and will be addressed in future releases:

DTaP/Tdap

- Scenario: DTaP Dose 5 in catch-up schedule forecasts with 6-month interval without regard to the individual's age.
- Scenario: Patient with no contraindication to Pertussis has Pediatric DT recorded in immunization history and does not meet definition of complete for Pertussis. Pediatric DT doses do not count towards series completion because it does not satisfy the Pertussis component. If patient has 5 doses in history, including Pediatric DT, and age at evaluation is < 7 yrs of age, a dose of DTaP should be forecast 28 days from the last dose.
- Tdap at 10 yrs of age marked as inadvertent dose but should satisfy adolescent booster dose.

HPV

Recommendation Change: HPV Forecasting for 26 years old, no history

- Scenario: CDSi 2016-0013 Female age 26, No HPV doses. Forecaster does not return a recommendation. Setting to suppress first dose is not enabled. Age indication expanded by CDC in October 2018.
- The ACIP recommendation was issued June 2019: **ACIP approved vaccination of persons age 27–45 years based on “shared clinical decision making” between the patient and clinician.** **Shared clinical decision making means the decision to vaccinate persons age 27 through 45 years should be based on a discussion of benefits and risks between the patient and the clinician.* This decision is not considered final until it is published in the MMWR. This ticket is blocked pending publication.

Recommendation Change: Harmonize HPV catch-up schedule for male and female

- In June 2019, ACIP voted unanimously to harmonize the routine catch-up vaccination schedule for both males and females through age 26. This decision is not considered final until it is published in the MMWR. This ticket is blocked pending publication.

Hib

Forecaster incorrectly recommends Dose #3 Hib at 4 week interval after dose 2, rather than an 8 week interval.

- Scenario: DOB 03/01/17 Dose #1 HIB-PRP-T on 05/19/17 at 2.6 months of age. Dose #2 was given on 10/05/17 at 7.2 months of age.
- Current behavior: Forecaster returned a recommended date of 11/02/20, only 4 weeks after the 2nd dose and the same as the minimum interval. This occurred because

forecasting for Hib was previously changed to forecast based on the last vaccination date/patient age to match CDSi.

- Expected behavior: If the forecasts evaluates the patient's current age as ≥ 12 months, the interval would be 8 weeks.
- Task: Determine method for forecasting based on supplied evaluation date. If evaluation date is null or "today", the forecast should be based on patient's current age.
- HDSAF-143

Forecaster correctly marks a Hib PRP-T dose as invalid but returns a "minimum interval not met" reason, rather than "minimum age not met".

- Scenario: DOB 08/23/2005. Patient received Hib (PRP-T) doses on 10/24/05, 01/10/2006, 02/27/2006, and 08/15/2006.
- Current behavior: Dose #4 is correctly marked as invalid. The reason for the invalid status displayed on the Vaccination Data Quality report is *Minimum interval from previous dose not met*.
- Expected behavior: Reason for invalid status is expected to display as *Minimum age for this dose not met*.
- HDSD-499

Recommended Hib intervals between Dose #1 and Dose #2 are different for Hib PRP-OMP and Hib-PRP-T and display "Minimum" interval dates as "Recommended" interval dates.

- Scenario: Forecaster returns recommended date for dose 2 of 4 weeks after dose 1 if Hib PRP-T Dose #1 is given at 3 mos.
- Current behavior: Forecaster returns recommended date for dose 2 of 8 weeks after dose 1 if Hib PRP-OMP Dose #1 is given at 3 mos.
- Expected behavior: Forecaster should return recommendation with 4 week interval when first dose is given before the 1st birthday. Forecaster should return recommendation with 8 week interval when first dose is given between 12-14 months.
- HDSD-567

Pneumococcal

Forecast returns PCV13 recommendation 1 year after inadvertent PPSV.

- Scenario: DOB 3/1/19, PPSV23 dose given 4/30/19.
- Current behavior: Forecast returns PCV13 recommendation 1 year after inadvertent PPSV. PPSV23 given at this age should not be considered to be part of the pneumococcal vaccination series. PCV13 should be administered as soon as the error is discovered.
- Expected behavior: Based on ACIP, PPSV23 given at this age should not be considered to be part of the pneumococcal vaccination series. PCV13 should be administered as soon as the error is discovered.
- HDSD-403, HDSD-455

Dose 2 incorrectly displays *Invalid PNEUMO (PCV): Minimum interval from previous dose not met.*

- Scenario #1: DOB: 08/10/2018, PCV13: 09/28/2018, 03/07/2019.
- Current behavior: Dose 2 incorrectly displays *Invalid PNEUMO (PCV): Minimum interval from previous dose not met.* Warning disappears with 4 day grace period.
- Expected behavior: Dose should not be marked as invalid.

- Scenario #2(related): DOB -8/30/2018, PCV 13 dose administered 03/28/2019
- Current behavior: Dose is incorrectly marked *Invalid PNEUMO (PCV): Minimum interval from previous dose not met,* even though it is the first dose on the record.
- Expected behavior: Dose should not be marked as invalid.
- HDS-422, HDS-418

Recommendation Change: PCV13 for Immunocompetent Older Adults

- The following change in recommendation for PCV13 in immunocompetent older adults was approved in June 2019: ACIP recommends PCV13 based on shared clinical decision making for adults 65 years and older who do not have an immunocompromising condition** and who have not previously received PCV13. All adults 65 years and older should receive a dose of PPSV23.*
- Of note, the recommendations for vaccination of adults at high risk of invasive pneumococcal disease (MMWR, Vol. 61, No. 40, pages 816-819) have not changed.
- This decision is not considered final until it is published in the MMWR. This ticket is blocked pending publication.

Rotavirus

Rotavirus Dose #2 dates off (Found by STC during regression testing)

- Scenario: CDSi Test Case 2013-0773 DOB: 05/17/2018, RV1 Dose #1 08/24/2018,
- Current behavior: Forecaster returns Min 9/21, rec 10/19, past due 11/18
- Expected behavior: Min and rec 9/21, past due 11/03

Two Rotavirus CDSI test cases with correct evaluation but inaccurate reason (Found by STC during regression testing)

Scenario #1: CDSI 2013-0782

- DOB 12/18/2018 , CVX 116 on 01/27/2019 , CVX 116 on 02/21/2019
- Current behavior: Minimum Interval from previous dose not met
- Expected behavior: Evaluation of Not Valid due to Minimum age for this dose not met.

Scenario #2 CDSI 2013-0785

- DOB 11/20/2018, CVX 116 on 12/29/2018 , CVX 116 on 01/26/2019, CVX 116 on 02/21/2019
- Current behavior: Minimum Interval from previous dose not met
- Expected behavior: Evaluation of Not Valid due to Minimum age for this dose not met.

Zostavax

Shingrix at age 18

- If a dose is inadvertently administered to an adult 18 through 49 years of age, CDC does not recommend repeating the dose but administering the second RZV dose on or after the 50th birthday. This guidance does not appear in the most recent zoster ACIP statement but is in the General Best Practices Guidance (Table 3-1 in the Timing and Spacing of Immunobiologics section at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/timing.html) and is based on guidance from CDC's zoster subject matter experts.
- Task: Determine approach to this documentation.
- HDS-821

Planned Logic Changes

New Vaccine Forecasting

- Addition for forecasting logic for Twinrix Jr.

Live Vaccine Rules

- HDS-535: Patient received OPV and MMR less than 28 days apart. MMR dose on 8/1/1988 is flagged as invalid but should be considered valid based on CDC General Recommendations on Immunization from the Pink Book. *"Parenteral live vaccines (MMR, MMRV, varicella, zoster, and yellow fever) and LAIV are not believed to have an effect on live vaccines given by the oral route (OPV, oral typhoid, and rotavirus). Live oral vaccines may be given at any time before or after live parenteral vaccines or LAIV."*
- HDS-519: Patient received RSV IGIV on 3/15/18 and then received MMR and varicella vaccines on 4/9/18. The MMR and varicella vaccines are incorrectly marked as invalid.

CDSi Logic NOT Being Implemented in ImmuCast 1.0

- CDSi allows a 5-dose Polio schedule when dose 4 is given too early (as in the use of combination vaccines). The 5-dose schedule considers an early Dose 4 as valid instead of invalid, similar to the Hepatitis B 4-dose schedule. ImmuCast 1.0 will not implement this logic due to the issues it will cause in the IWeb school certificates and reports.

Product Documentation

Product documentation is located on the STC Documentation Portal:
<https://documentation.stchome.com/>.

The following documents are available for this version of ImmuCast:

- ImmuCast 5.18.1 User Guide
- ImmuCast 5.24.0 Release Notes